


BULLETIN OF AMERICAN ODONATOLOGY



Volume 8 Number 1

5 Apr 2004

**DISTRIBUTION OF NORTH AMERICAN ODONATA. PART II:
MACROMIIDAE, CORDULIIDAE AND LIBELLULIDAE**

Thomas W. Donnelly, p. 61 - 32

THE DRAGONFLY SOCIETY OF THE AMERICAS

Business address: c/o T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903

EXECUTIVE COUNCIL 2003 - 2005

| | | |
|-------------------------------|---------------|------------------|
| President | R. Beckemeyer | Wichita KS |
| President Elect | S. Krotzer | Centreville AL |
| Immediate past President | D. Paulson | Seattle WA |
| Vice President, Canada | R. Cannings | Victoria, BC |
| Vice President, Latin America | R. Novelo G. | Jalapa, Veracruz |
| Secretary | S. Dunkle | Plano TX |
| Treasurer | J. Daigle | Tallahassee FL |
| Editor | T. Donnelly | Binghamton NY |
| Regular member | J. Abbott | Austin TX |
| Regular member | S. Valley | Albany OR |
| Regular member | S. Hummel | Lake View IA |

JOURNALS PUBLISHED BY THE SOCIETY

ARGIA, the quarterly news journal of the **DSA**, is devoted to non-technical papers and news items relating to nearly every aspect of the study of Odonata and the people who are interested in them. The editor especially welcomes reports of studies in progress, news of forthcoming meetings, commentaries on species, habitat conservation, noteworthy occurrences, personal news items, accounts of meetings and collecting trips, and reviews of technical and non-technical publications. Articles for publication in **ARGIA** should preferably be submitted as hard copy and (if over 500 words) also on floppy disk (3.5" or 5.25"). The editor prefers MS DOS based files, preferably written in WORD, WORD for WINDOWS, WordPerfect, or WordStar. Macintosh WORD disks can be handled. All files should be submitted **unformatted and without paragraph indents**. Each submission should be accompanied by a text (=ASCII) file. Other languages should be submitted only as text (=ASCII) files. Line drawings are acceptable as illustrations.

T. Donnelly (address below) is the interim editor of **ARGIA**.

BULLETIN OF AMERICAN ODONATOLOGY is devoted to studies of Odonata of the New World. This journal considers a wide range of topics for publication, including faunal synopses, behavioral studies, ecological studies, etc. The **BAO** publishes taxonomic studies but will not consider the publication of new names at any taxonomic level. Enquiries and submission of manuscripts should be made to **BAO** editor T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903. Final submissions (after review) should be made on floppy disk, as above, with illustrations in final form and preferably adjusted to final size.

MEMBERSHIP IN THE DRAGONFLY SOCIETY OF THE AMERICAS

Membership in the **DSA** is open to any person in any country. Dues for individuals in the US, Canada, or Latin America are \$15 for regular membership and \$20 for institutions or contributing membership, payable annually on or before 1 March of membership year. Dues for members in the Old World are \$25.

Dues should be mailed to Jerrell Daigle, 2067 Little River Lane, TALLAHASSEE FL 32311

The **BULLETIN OF AMERICAN ODONATOLOGY** is available by a separate subscription at \$15 for members and \$18.75 for non-members and institutions.

DISTRIBUTION OF NORTH AMERICAN ODONATA. PART II: MACROMIIDAE,
CORDULIIDAE, LIBELLULIDAE¹

Thomas W. Donnelly, 2091 Partridge Lane, Binghamton NY 13903, tdonelly@binghamton.edu

ABSTRACT: Dot-map presentations show the distributions for species of Macromiidae, Corduliidae, and Libellulidae in North America. Additional comments, including subspecies information, are provided for several species.

INTRODUCTION: This is the second of a series that started with Donnelly (2004a). Introductory and explanatory comments in that number will not be repeated here.

SUBSPECIES, AND SPECIES OF SPECIAL INTEREST:

Macromia illinoensis: Donnelly and Tennesen (1994) described structural variations in the taxa *georgina* and *illinoensis* and proposed that they had a subspecific relationship. They noted four counties where the taxa varied smoothly between the subspecific end members, and noted that in a fifth county (Macon Co. NC) there were separate population of the two which did not so vary. May and Carle (1996) noted that "The two taxa seem distinct in structural characters in New Jersey and are sympatric in Virginia and North Carolina." We see no reason to alter our conclusion.

Macromia X wabashensis: I treat this as a hybrid, probably between *taeniolata* and *pacifica*. The Texas specimens might alternatively have *annulata* as one parent.

Epithea costalis, cynosura, and petechialis: These taxa have caused extensive difficulties. Recent examination of numerous specimens from eastern Texas, Oklahoma, Kansas, and Colorado, have shown unspotted *petechialis* is far more widespread than previously recognized. Probably plains (western Kansas, Colorado, western Oklahoma, central and western Texas) records of "*costalis*" and "*cynosura*" are probably pale-winged *petechialis*. The only reliable method of distinguishing among these three taxa is by a close examination of the male cerci and abdominal base.

In several localities in Ohio and scattered localities elsewhere, *costalis* and *cynosura* appear to intergrade freely, although in other localities, no evidence of intergradation has been detected. While these species appear to be valid biological

species, the extent of, and the reasons for, their apparent intergradation are not yet understood.

Epithea semiaquea: There is a broad gap in the distribution of this species, with no records from north-central Florida to Texas. Specimens from Massachusetts appear to lack the wing spots and have been called *cynosura*.

Somatochlora hineaea: The records from Alabama, Ohio, and Indiana are historic.

Williamsonia lintneri: The record from Albany, New York is historic.

Erythemis simplicicollis and collocata: The apparent subspecific relationship between these two taxa was discussed by Donnelly (2004b).

Leucorrhinia proxima: The eastern and western forms are different in both appearance and habits. The eastern forms have yellow pale color and become quite pruinose. They commonly do not perch in conspicuous positions. The western forms are patterned with bright red, are far less pruinose, and perch in very exposed positions. The boundary between the two forms appears to be approximately in western Wisconsin.

Orthemis antillean species: This species has not been named. It flies with ferruginea around Miami, and the two can be distinguished by several differences in appearance. The antillean form ranges to Trinidad, but is not certainly specifically distinct with a co-occurring, and also unnamed, purple form in Hispaniola, Puerto Rico, and the Virgin Islands. It is premature to name any of these forms.

Perithemis tenera: This species appears to grade smoothly into the tropical "species" *mooma*, and several specimens from southern Texas could be referred to *mooma*. It is more realistic to regard them as one wide ranging and variable species. Some of the Mexican records shown are of "*mooma*", but Mexican records of *tenera* are numerous and widespread

Platthemis lydia: Eastern and western forms, distinguished by the shape of the styli on the basal abdominal segment (Donnelly, 1991), grade

¹Bulletin of American Odonatology 8(1) 1:32

together in a narrow zone in west-central Montana to western Wyoming.

Sympetrum internum and *rubicundulum*: Throughout most of their considerable overlapping ranges, these appear to be easily distinguishable species (Williamson, 1933). Along the Atlantic seaboard, from at least New Jersey to Cape Cod, the most abundant form appear to be *rubicundulum*, but with a hamule intermediate between this species and *internum*. Specimens of apparently valid *internum* are found sporadically in the same area.

Sympetrum internum and *obtrusum*: Hybrids between these two species have been found at several northeastern localities (Donnelly, 1991).

Sympetrum semicinctorum: Specimens taken in westernmost Ontario and Wisconsin, and in Northeastern Minnesota, appear to be transitional between *semicinctorum* and *occidentale*. I believe the two to have a subspecific relation, as shown. I have not further differentiated californicum and *fasciatum*.

REFERENCES CITED

Donnelly, T.W., 1991, Current problems in *Sympetrum* and *Libellula* (a.k.a. *Plathemis*). ARGIA 3(4): 8-12


Donnelly, T.W., 2004a, Distribution of North American Odonata. Part I: Aeshnidae, Petaluridae, Gomphidae, and Cordulegastridae. Bulletin of American Odonatology 7(4): 61-90

Donnelly, T.W., 2004b, *Erythemis simplicicollis* and *collocata* – subspecies? ARGIA 15(4): 11-13

Donnelly, T.W., and Tennessen, K.J., 1994, *Macromia illinoensis* and *georgina*: a study of their variation and apparent subspecific relationship (Odonata: Corduliidae). Bulletin of American Odonatology 2(3): 27-61

May, M.L., and Carle, F.L., 1996, An annotated list of the Odonata of New Jersey. Bulletin of American Odonatology, 4(1): 1-35

Williamson, E.B., 1933, The status of *Sympetrum assimilatatum* (Uhler) and *Sympetrum decisum* (Hagen) (Odonata – Libellulinae). Occ. Papers Mus. Zoology. Univ. Michigan, no. 264, 7 p, 1 pl.



INDEX TO MAPS

(NOTE: M refers to Macromiidae, C to Corduliidae, and L to Libellulidae)

- Didymops floridensis M1
- Didymops transversa M1
- Macromia alleghaniensis M2 C1
- Macromia annulata M1
- Macromia illinoensis M2 C1
- Macromia magnifica M1
- Macromia margarita M1
- Macromia pacifica M1
- Macromia X wabashensis M2 C1
- Macromia taeniolata M1
- Cordulia shurtleffi M2 C1
- Dorocordulia lepida M2 C1
- Dorocordulia libera M2 C1
- Epitheca (Epicordulia) princeps C3
- Epitheca (Tetragoneuria) canis C3
- Epitheca (Tetragoneuria) costalis C2
- Epitheca (Tetragoneuria) cynosura C2
- Epitheca (Tetragoneuria) petechialis C2
- Epitheca (Tetragoneuria) semiaquea C2

- Epitheca (Tetragoneuria) sepia C2
- Epitheca (Tetragoneuria) spinigera C2
- Epitheca (Tetragoneuria) spinosa C2
- Epitheca (Tetragoneuria) stella C2
- Helocordulia selysii M2 C1
- Helocordulia uhleri C4
- Neurocordulia alabamensis C3
- Neurocordulia michaeli C3
- Neurocordulia molesta C4
- Neurocordulia obsoleta C4
- Neurocordulia virginiana C3
- Neurocordulia xanthosoma C3
- Neurocordulia yamaskanensis C4
- Somatochlora albicincta C5
- Somatochlora brevicincta C6
- Somatochlora calverti C7
- Somatochlora cingulata C5
- Somatochlora elongata C8
- Somatochlora ensigera C8
- Somatochlora filosa C8
- Somatochlora forcipata C7
- Somatochlora franklinII C7
- Somatochlora georgiana C8
- Somatochlora hineana C4








| | |
|--|--|
| <i>Somatochlora hudsonica</i> C6 | <i>Leucorrhinia patricia</i> L9 |
| <i>Somatochlora incurvata</i> C10 L1 | <i>Leucorrhinia proxima</i> L8 |
| <i>Somatochlora kennedyi</i> C7 | <i>Libellula auripennis</i> L10 |
| <i>Somatochlora linearis</i> C8 | <i>Libellula axilena</i> L10 |
| <i>Somatochlora margarita</i> C7 | <i>Libellula comanche</i> L10 |
| <i>Somatochlora minor</i> C9 | <i>Libellula composita</i> L11 |
| <i>Somatochlora ozarkensis</i> C8 | <i>Libellula croceipennis</i> L10 |
| <i>Somatochlora provocans</i> C9 | <i>Libellula cyanea</i> L10 |
| <i>Somatochlora sahlbergi</i> C6 | <i>Libellula flavida</i> L9 |
| <i>Somatochlora semicircularis</i> C6 | <i>Libellula forensis</i> L8 |
| <i>Somatochlora septentrionalis</i> C6 | <i>Libellula incesta</i> L11 |
| <i>Somatochlora tenebrosa</i> C9 | <i>Libellula jesseana</i> L9 |
| <i>Somatochlora walshii</i> C9 | <i>Libellula luctuosa</i> L12 |
| <i>Somatochlora whitehousei</i> C5 | <i>Libellula needhami</i> L10 |
| <i>Somatochlora williamsoni</i> C5 | <i>Libellula nodisticta</i> L9 |
| <i>Williamsonia fletcheri</i> C4 | <i>Libellula pulchella</i> L12 |
| <i>Williamsonia lintneri</i> C4 | <i>Libellula quadrimaculata</i> L12 |
| <i>Brachymesia furcata</i> C10 L1 | <i>Libellula saturata</i> L11 |
| <i>Brachymesia gravida</i> L2 | <i>Libellula semifasciata</i> L11 |
| <i>Brachymesia herbida</i> C10 L1 | <i>Libellula vibrans</i> L11 |
| <i>Brechmorhoga mendax</i> C10 L1 | <i>Macrodiplax balteata</i> L2 |
| <i>Brechmorhoga pertinax</i> C10 L1 | <i>Macrothemis imitans</i> L2 |
| <i>Cannaphila insularis</i> C10 L1 | <i>Macrothemis inacuta</i> C10 L1 |
| <i>Celithemis amanda</i> L5 | <i>Macrothemis inequiunguis</i> L2 |
| <i>Celithemis berthia</i> L5 | <i>Miathyria marcella</i> L5 |
| <i>Celithemis elisa</i> L7 | <i>Micrathyria aequalis</i> L3 |
| <i>Celithemis eponina</i> L7 | <i>Micrathyria didyma</i> L3 |
| <i>Celithemis fasciata</i> L7 | <i>Micrathyria hagenii</i> L3 |
| <i>Celithemis martha</i> L6 | <i>Nannothemis bella</i> L11 |
| <i>Celithemis ornata</i> L7 | <i>Orthemis antillean species</i> L13 |
| <i>Celithemis verna</i> L6 | <i>Orthemis discolor</i> L13 |
| <i>Crocothemis servilia</i> C10 L1 | <i>Orthemis ferruginea</i> L13 |
| <i>Dythemis fugax</i> L2 | <i>Pachydiplax longipennis</i> L14 |
| <i>Dythemis maya</i> L2 | <i>Paltothemis lineatipes</i> L3 |
| <i>Dythemis nigrescens</i> L2 | <i>Pantala flavescens</i> L14 |
| <i>Dythemis velox</i> L2 | <i>Pantala hymenaea</i> L14 |
| <i>Erythemis attala</i> L4 | <i>Perithemis domitia</i> L13 |
| <i>Erythemis peruviana</i> L4 | <i>Perithemis intensa</i> L13 |
| <i>Erythemis plebeja</i> L5 | <i>Perithemis tenera</i> L13 |
| <i>Erythemis simplicicollis</i> L4 | <i>Plathemis lydia</i> L15 |
| <i>Erythemis vesiculosa</i> L5 | <i>Plathemis subornata</i> L15 |
| <i>Erythrodiplax basifusca</i> L4 | <i>Pseudoleon superbus</i> L5 |
| <i>Erythrodiplax berenice</i> L4 | <i>Sympetrum ambiguum</i> L15 |
| <i>Erythrodiplax funerea</i> L3 | <i>Sympetrum corruptum</i> L17 |
| <i>Erythrodiplax fusca</i> L4 | <i>Sympetrum costiferum</i> L16 |
| <i>Erythrodiplax minuscula</i> L3 | <i>Sympetrum danae</i> L16 |
| <i>Erythrodiplax umbrata</i> L5 | <i>Sympetrum illotum</i> L8 |
| <i>Idiataphe cubensis</i> L3 | <i>Sympetrum internum</i> L17 |
| <i>Ladona deplanata</i> L6 | <i>Sympetrum madidum</i> L15 |
| <i>Ladona exusta</i> L6 | <i>Sympetrum obtrusum</i> L17 |
| <i>Ladona julia</i> L7 | <i>Sympetrum pallipes</i> L15 |
| <i>Leucorrhinia borealis</i> L9 | <i>Sympetrum rubicundulum</i> L15 |
| <i>Leucorrhinia frigida</i> L7 | <i>Sympetrum semicinctum</i> L16 |
| <i>Leucorrhinia glacialis</i> L8 | <i>Sympetrum signiferum</i> L13 |
| <i>Leucorrhinia hudsonica</i> L8 | <i>Sympetrum vicinum</i> L18 |
| <i>Leucorrhinia intacta</i> L9 | <i>Sympetrum internum X obtrusum</i> L17 |

Tauriphila australis L4
Tauriphila azteca L4
Tholymis citrina L3
Tramea abdominalis L6
Tramea binotata L5

Tramea calverti L6
Tramea carolina L6
Tramea insularis L5
Tramea lacerata L18
Tramea onusta L18







- ◆ Adult specimen; often used for stage unspecified
- Photograph of adult
- ▮ Larva or exuviae
- ◇ Sight Record
- Record specified for state (US or Mexico) or region

Key to the symbols used in the maps.




MACROMIIDAE 1


Didymops floridensis, transversa
Macromia pacifica, annulata, georgina, magnifica, taeniolata
margarita






MACROMIIDAE 2; CORDULIIDAE 1

Macromia illinoensis, allegheniensis, X wabashensis
Cordulia shurtleffi
Dorocordulia libera, lepida
Helocordulia selysii















Many western *cynosura* records may be clear-winged *petechialis*.



Many western *costalis* records may be clear-winged *petechialis*.










CORDULIIDAE 2 *Epitheca cynosura, costalis, petechialis, sepia, stella, spinosa*



CORDULIIDAE 3




Epitheca canis, spinigera, princeps

Neurocordulia alabamensis, michaeli, virginiensis, xanthosoma









CORDULIIDAE 4






- Helocordulia uhleri*
- Neurocordulia yamaskanensis, molesta, obsoleta*
- Williamsonia fletcheri, lintneri*
- Somatochlora hineana*









CORDULIIDAE 5 *Somatochlora albicincta, cingulata, whitehousei*







CORDULIIDAE 6 *Somatochlora brevicincta, sahlbergi, septentrionalis, hudsonica, semicircularis, williamsoni*











CORDULIIDAE 7 *Somatochlora kennedyi, calverti, margarita, forcipata, franklinii*



CORDULIIDAE 8 *Somatochlora elongata, ensigera, filosa, linearis, georgiana, ozarkensis*











CORDULIIDAE 9 *Somatochlora provocans, tenebrosa, minor, walshii*










CORDULIIDAE 10; LIBELLULIDAE 1

- Somatochlora incurvata*
- Brachymesia furcata*, *herbida*
- Brechmorhoga mendax*, *pertinax*
- Macrothemis inacuta*
- Cannaphila insularis*










LIBELLULIDAE 2

- Brachymesia gravida*
- Macrothemis imitans, inequiunguis*
- Macrodiplax balteata*
- Dythemis fugax, maya, nigrescens, velox*




LIBELLULIDAE 3

- Idiataphe cubensis*
Paltothermis lineatipes
Micrathyria aequalis, didyma, hagenii
Erythrodiplax funerea, minuscula
Tholymis citrina




LIBELLULIDAE 4


Erythemis simplicicollis, attala, peruviana
Erythrodiplax basifusca, fusca
Tauriphila australis, azteca




Erythrodiplax umbrata




Erythemis vesiculosa




Erythemis plebeja



Miathyria marcella




Tramea insularis




Pseudoleon superbus



Tramea binotata










Celithemis amanda









Celithemis bertha

- LIBELLULIDAE 5** *Erythrodiplax umbrata*
Erythemis vesiculosa, plebeja
Tramea insularis, binotata
Pseudoleon superbus
Miathyria marcella
Celithemis amanda, bertha








LIBELLULIDAE 6

Ladona deplanata, exusta
Tramea calverti, carolina,
abdominalis
Celithemis verna, martha









LIBELLULIDAE 7

- Celithemis elisa, ornata, eponina, fasciata*
Celithemis ornata
Leucorrhinia frigida









LIBELLULIDAE 8

Leucorrhinia proxima, glacialis, hudsonica
Libellula forensis
Sympetrum illotum









LIBELLULIDAE 9




Leucorrhinia patricia, intacta, borealis
Libellula nodisticta, flavida, jesseana










LIBELLULIDAE 10 *Libellula auripennis, axilena, comanche, croceipennis, cyanea, needhami*



LIBELLULIDAE 11
Libellula incesta, vibrans, composita,
saturata, semifasciata
Nannothemis bella






LIBELLULIDAE 12 *Libellula pulchella, luctuosa, quadrimaculata*








LIBELLULIDAE 13

Orthemis ferruginea, discolor, antillean species
Perithemis domitia, intensa, tenera
Sympetrum signiferum






LIBELLULIDAE 14 *Pachydiplax longipennis*; *Pantala flavescens, hymenaea*







LIBELLULIDAE 15




Plathemis lydia, subornata
Sympetrum pallipes, madidum, ambiguum, rubicundulum



LIBELLULIDAE 16 *Sympetrum semicinctum, danae, costiferum*



LIBELLULIDAE 17 *Sympetrum corruptum, internum, obtusum (incl. hybrid)*



LIBELLULIDAE 18 *Tramea lacerata, onusta*
Sympetrum vicinum