

ARGIA

THE NEWS JOURNAL OF THE DRAGONFLY SOCIETY OF THE AMERICAS

VOLUME 16

5 April 2004

NUMBER 1



PUBLISHED BY THE DRAGONFLY SOCIETY OF THE AMERICAS

THE DRAGONFLY SOCIETY OF THE AMERICAS

Business address: c/o T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903

EXECUTIVE COUNCIL 2003 - 2005

President	R. Beckemeyer	Wichita KS
President Elect	S. Krotzer	Centreville AL
Immediate past President	D. Paulson	Seattle WA
Vice President, Canada	R. Cannings	Victoria, BC
Vice President, Latin America	R. Novelo G.	Jalapa, Veracruz
Secretary	S. Dunkle	Plano TX
Treasurer	J. Daigle	Tallahassee FL
Editor	T. Donnelly	Binghamton NY
Regular member	J. Abbott	Austin TX
Regular member	S. Valley	Albany OR
Regular member	S. Hummel	Lake View IA

JOURNALS PUBLISHED BY THE SOCIETY

ARGIA, the quarterly news journal of the **DSA**, is devoted to non-technical papers and news items relating to nearly every aspect of the study of Odonata and the people who are interested in them. The editor especially welcomes reports of studies in progress, news of forthcoming meetings, commentaries on species, habitat conservation, noteworthy occurrences, personal news items, accounts of meetings and collecting trips, and reviews of technical and non-technical publications. Articles for publication in **ARGIA** should preferably be submitted as hard copy and (if over 500 words) also on floppy disk (3.5" or 5.25"). The editor prefers Windows files, preferably written in WORD, WORD for WINDOWS, WordPerfect, or WordStar. Macintosh WORD disks can be handled. **ALL FILES SHOULD BE SUBMITTED UNFORMATTED AND WITHOUT PARAGRAPH INDENTS.** Each submission should be accompanied by a text (=ASCII) file. Other languages should be submitted only as text (=ASCII) files. Line drawings are acceptable as illustrations.

T. Donnelly (address below) is the interim editor of **ARGIA**.

BULLETIN OF AMERICAN ODONATOLOGY is devoted to studies of Odonata of the New World. This journal considers a wide range of topics for publication, including faunal synopses, behavioral studies, ecological studies, etc. The **BAO** publishes taxonomic studies but will not consider the publication of new names at any taxonomic level. Enquiries and submission of manuscripts should be made to **BAO** editor T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903. Final submissions (after review) should be made on floppy disk, or as e-mail attachment, as above, with illustrations in final form and preferably adjusted to final size.

MEMBERSHIP IN THE DRAGONFLY SOCIETY OF THE AMERICAS

Membership in the **DSA** is open to any person in any country. Dues for individuals in the US, Canada, or Latin America are \$15 for regular membership and \$20 for institutions or contributing membership, payable annually on or before 1 March of membership year. Dues for members in the Old World are \$25.

Dues should be mailed to Jerrell Daigle, 2067 Little River Lane, TALLAHASSEE FL 32311

The **BULLETIN OF AMERICAN ODONATOLOGY** is available by a separate subscription at \$15 for members and \$18.75 for non-members and institutions.

Front cover: *Gomphus oklahomensis*, San Jacinto Co. TX. Photo by Greg Lasley

In This Issue

Blowing snow in April leads me to desolation. My friends are joyously e-mailing me their results from the sunny south. Greg Lasley sent by e-mail a beautiful photo (which is the cover photo) of *Gomphus oklahomensis* from a lovely pond in east Texas that I directed him to a week ago. Looking at the cover reminds me of how I enjoyed walking around this pond when we lived in Houston. I can't tell you how much I wish I were there right now!

Meetings are just ahead of us. The Eglin AFB meeting will be under way even as you read this, but the SE meeting in Kentucky is in June. News of this year's Eagle Hill seminar did not reach me in time for the last issue, but here it is now.

Michael Veit tells us that his Nymph Fest was a great success. This tells us that even in the depressing depths of winter there are useful and fun things to do in the world of bugs.

Roy Beckemeyer sends us installment 2 of his series of articles on insect flight. It is about time that someone filled us in on how these amazing bugs do it. It is a challenging story, but well worth the effort, especially to those who opted out of physical science in their youth.

The sense of macabre whimsy that April snow storms engendered in me caused me to place immediately after Roy's article the text of an ad on the web for a flying dragonfly toy from Korea. Do you remember the famous line from Monty Python, "And now for something completely different!"?

And now indeed for something completely different. Paul Catling tells us that the Northwest Territories, which have even more winter than Binghamton, also have lots of great dragonflies. If you can protect yourself from the biting flies, then this is the place to be. At least for a few weeks in the middle of the summer.

The text of an article on a lawsuit regarding habitat for *Somatochlora hineana* (aka Hine's Emerald), our only federally endangered dragonfly will be of interest for the conservation minded.

It's Roy again with a thought-provoking discussion of a recent article in *NATURE* on the origins of the insects. One of the amazing results of Paleozoic paleontology is the speed with which the land

surface was populated by rapidly evolving plants and animals following the first appearance of terrestrial algae-like organisms.

Jerrell, spraying exclamation marks like pepper spray, tells us of an apparently jolly and certainly productive trip to south Florida with Ken Tennessen. This is another great way to beat the winter.

Kent Fothergill and friends report *Cordulegaster dorsalis* from Idaho, which is another state that is undersurveyed. Perhaps in a few years the lists from many parts of the west and great plains will start filling up. This is a good start.

A state which needs no apologies is Iowa. Steve Hummel shows us that even in one of the best-known states it is possible to extend the list. In the same vein, Dwayne Sabine and friends extend the impressive list for New Brunswick even more.


Moving south, François Meurgey sends in three articles – one on Guadeloupe and one on the Dominican Republic (both of which are a lot warmer in the winter than Binghamton). His third contribution is on our old friend *Anax*, which has recently showed up in France. If you see a Green Darner carrying a teeny-tiny baguette, you will know where it spent the winter.

We have a lot to review this time. Allen Barlow reviews Dick Walton's recent DVD on eastern dragonflies, which is the successor of his earlier video on the same subject. This is a must for beginners. Similarly, Cynthia Berger's "Dragonflies" belongs on every shelf. I also review an attractive children's book (there are far too few children's books available) will be of wide interest, especially to grandparents.

My, how times change. Last year I didn't even know what Powerpoint was. Now I am reviewing a Powerpoint presentation, by May Lenzer.

Roy Beckemeyer, who always seems to pick up interesting things that others miss, reviews a website on egg parasitism. Maybe you can really learn everything on the web.

Some breaking news at the end describes a recent irruption of *Paltothemis lineatipes* (Red Rock Skimmer) in southern California.



Calendar: 2004 DSA and other meetings			
meeting	date	place	contact
Eglin AFB Mtg.	9-11 Apr	Niceville FL	jdaigle@nettally.com
New England Conf.	17 Apr	Athol MA	odenews@odenews.net
Dragonfly Days	10-14 May	Weslaco TX	http://www.valleynaturecenter.org
			http://www.valleynaturecenter.org
Eagle Hill seminar	30 May - 5 June	Eagle Hill, ME	office@eaglehill.us
DSA SE Mtg.	10-13 June	Mammoth Cave KY	bugman@scrtc.com
GLOM	22-26 June	Cleveland OH	rboronka@cmnh.org
DSA NE Mtg.	24-27 June	Hartford VT	bryan@vermontbirdtours.com
			bryan@vermontbirdtours.com
DSA National Mtg.	23-24 Aug	Decorah IA	mshummel@netins.net
Dragonfly Festival	28-29 Aug	Roswell NM	http://www.rt66/~kjherman/fbl/df.html

ANNOUNCING: THE 2004 ADVANCED AND PROFESSIONAL LEVEL ENTOMOLOGY SEMINARS AT THE HUMBOLDT INSTITUTE ON THE COAST OF EASTERN MAINE!

Damselflies and Dragonflies: Systematics and Biomonitoring, May 30 - June 5

This seminar is offered for naturalists and specialists involved with or interested in aquatic insect inventory and bio-monitoring. The program will cover several topics in the study of Odonata including their life cycle, morphology, behavior, and species distributions in the northeast. Sampling and preservation techniques, and both printed and electronic (WWW) references will be reviewed. Fieldwork will provide experience in survey methods for all life stages; larvae and exuviae will be sampled for taxonomic study and photography. Lab work will focus on identification, as well as the use and creation of virtual collections using digital images. Reference specimens will be provided, but additional material is welcome. Frederick H. SaintOurs (fred.saintours@umb.edu) has been documenting and studying Odonata throughout the northeast and the American tropics for the last seven years. He has worked as consultant in rare species surveys, and has been contributing records for state-wide inventory programs throughout New England as well as articles for Ode News and the New England Chapter of the Society for Conservation Biology. He has organized and led numerous surveys and workshops on aquatic macroinvertebrate monitoring and identification in streams and rivers through grants received from the

National Science Foundation, Massachusetts Environmental Trust, Natural Heritage and Endangered Species Program, and Riverways Program. Currently he is developing image-based, web-accessible invertebrate identification tools as part of the Electronic Field Guide (EFG) project at U-Mass Boston. Other advanced entomology seminars ...

June 6 - 12 Macrolepidoptera: Identification and Systematics ... Brian Scholtens

June 20 - 26 ... The EPT Taxa: Systematics and Biomonitoring: Ephemeroptera, Plecoptera, and Trichoptera ... Steven Burian Jun 26 - Jul 2 . Beetles ... Donald Chandler

July 11 - 17 ... Chironomids: Systematics and Biomonitoring ... Leonard Ferrington

July 11 - 17 ... Photography Through the Microscope and Close-up Photography ... Stanley Cichowitz

For more information, please contact Humboldt Institute, PO Box 9, Steuben, ME 04680-0009.

207-546-2821. Fax 207-546-3042.

E-mail: <office@eaglehill.us>

Online registration and information
<http://www.eaglehill.us>

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

**SOUTHEAST DSA MEETING
UPDATED INFORMATION**

Carl Cook

It's that time! Let's all get our plans finalized to attend DSA's Southeast Regional Meeting at

Best Western, (270) 773-3161, (toll free 800-528-1234), \$65.00 tax included
Howard Johnson, (270) 773-2181, (toll free 800-321-4245), \$65.00 tax included
Ramada Limited, (270) 773-3121, (no toll free) \$62.00 tax included
Days Inn, (270) 773-2151, (no toll free) \$64.00 tax included
Quality Inn, (270) 773-3101, (no toll free) \$54.00 tax included


The operations center for the meeting is being provided by the Science Management Division of MCNP, it is known as Hamilton Valley Research Center. This facility has just about every contrivance we could possibly need for the meeting, a large conference room for presentations (bring those slides!), lab rooms for specimen preparation, and ten rooms for over-night housing, everything is air conditioned. The rooms, however, are unfurnished- bring your own bedrolls or bags- the rooms will hold two persons and cost \$4.00 per person per night. We will assemble at the Center each morning at 8:00 AM CDT to begin daytime activities and again following dinner for evening programs.

would be glad to hear from everyone planning to attend, or to answer your questions at (270) 565-3795 or <bugman@scrtc.com>


You still have time to sign up for this one-day meeting, co-sponsored by the Massachusetts Natural Heritage & Endangered Species Program, the Athol Bird and Nature Club, and Ode News. Speakers will explore distribution, conservation and

edge from the problem of how the pressure varies from the base of the wing to the tip.

Let's begin by looking at how the pressure is distributed along the chord of the wing – the leading to trailing edge (Figure 2). We will




be looking at the wing end-on, from the wing tip, at a representative location about midway out on the wing. We will represent the wing cross section as a straight line to begin with, and will use more representative cross sections later.



We will show the air movement past the wing by using what are called streamlines. Streamlines are lines which are always parallel to the air velocity.

This means that there is no air flowing across the streamlines. We could demonstrate streamlines in a wind tunnel by injecting smoke into the airstream through evenly spaced tubes so that the smoke enters parallel to the flowing air. In the steady




flow of the tunnel, the lines formed by the smoke would be equivalent to streamlines.

Figure 3 shows the streamlines for air flowing past a circular cylinder. The streamlines are parallel and horizontal far to the left of the cylinder, representing a horizontal uniform flow of air in the tunnel. The streamlines are once again parallel and horizontal far to the right. However, they curve around the cylinder as they approach it, and the streamlines get farther apart just before they reach the cylinder and as they begin to go around it. They get closer together as they pass by the top and bottom of the cylinder. Then they spread apart again, finally ending up parallel once the air is far enough beyond the cylinder.

The air flowing along or near the line passing through the center of the cylinder slows down because the cylinder blocks the flow. As the air flows up or down and around the cylinder, it gets squeezed into a smaller area and accelerates. The flow speed is depicted in Fig. 3 by the arrows. Far to the left they are all the same length. At the top and bottom of the cylinder, the arrows are longer. Far to the right they are the same length again.

Since no air flows across the streamlines, the mass of air per second passing through any point between two streamlines must be the same. Since



the air at these low speeds has constant density, that means that when streamlines get closer together, the air passing between them must be flowing faster than where they are farther apart.


You may recall Bernoulli's Equation from your high school physics class:

$\text{PRESSURE} \times \frac{1}{2} \text{VELOCITY}^2 = \text{CONSTANT}$. It basically states that the energy of the flow between any two streamlines is conserved. It also says that, wherever two streamlines get further apart, the velocity goes down and therefore the pressure goes up and wherever two streamlines get closer together, the velocity goes up and the pressure goes down.

The bottom half of Figure 3 shows the pressure distribution over the upper and lower surfaces of the cylinder resulting from the flow pattern shown in the top half of the figure. In this case the pressure is given in psi and a horizontal line at 14.7 psi marks atmospheric pressure. Since the flow is symmetrical, the pressure is the same on the upper and lower halves of the cylinder, and the curves overlap. The curve shows that the pressure near the front and back sides of the cylinder is greater than atmospheric, and that over the rest of the surface is less than atmospheric. Since the pressure is


symmetrical, there is no net force acting on the cylinder.

Figure 4 shows the flow past a horizontal flat plate. When the plate is horizontal, there is no differential




pressure generated and the pressure acting on the plate is simply the atmospheric pressure (in this and the rest of the figures, the pressure is in metric units – atmospheric pressure is 101.26 kilo-Pascals). In order to generate aerodynamic lift, we need to either alter the shape of our wing or incline it at an angle to the flow.

Figure 5 shows the flow past our plate when it is inclined at a positive angle of attack of 15 degrees. Note how the streamlines beneath the wing widen (flow decelerates and pressure increases) and those above the wing get closer together (flow accelerates and pressure decreases). The narrowest region is that of the streamline that curves sharply around the leading edge of the wing. Look at the pressure plot. The solid curve is the pressure on the upper surface of the wing, the broken curve the pressure on the lower side. Note that the pressure on the upper side is less than atmospheric pressure, and it becomes very much less near the leading edge. The pressure on the lower surface is greater than atmospheric for most of the wing chord, becoming less than atmospheric pressure in the small area just below the leading edge. The combination of higher pressure below the wing and lower above the wing yields a net supporting pressure that equals the lift




of the wing. That very large lower than atmospheric pressure dip on the upper wing surface at near the leading edge is called "leading edge suction", and occurs on all wings. It is the need for the flow to accelerate around the leading edge that eventually causes the wing to "stall" and lose its lift at very high angles of attack (more on this in a later installment).



We can make a wing in a shape that will generate lift without inclining it at an angle to the flow. Figure 6 shows a flat plate that has been curved to have a convex upper surface (called positive "camber"). Again we see wide spacing of lower surface streamlines and narrow spacing of upper surface ones. This yields a greater than

atmospheric pressure on the lower surface and a



lower pressure on the upper surface and therefore a net lift force.

In Figure 7 this cambered wing is inclined at 15 degrees. Compare this flow field and pressure distribution to that of Fig. 5 and you will see that the cambered wing generates higher lift than the flat wing at the same angle of attack. We see that it would be beneficial for insects to have wings that curve into a cambered shape under load rather than to stay flat.

It is usually customary to subtract the pressure on the upper wing surface from the pressure on the lower wing surface and to show the net pressure distribution on the wing as we have done in Figure 8 for a flat plate wing at 15 degrees inclination. We still see that much of the lift on the wing is generated by large pressure at the leading edge – certainly not a uniform or constant pressure – but note that the average pressure still needs to add up to the wing loading – 5 N/sq m for our *Anax junius*.

The final picture, Figure 9, shows the pressure distribution over a wing having the cross sectional shape of an *Anax junius* hind wing. You can see the corrugations at the leading edge (left end). Those corrugations, of course, stiffen and strengthen the wing so that it resists bending under pressure. And, amazingly enough, they are concentrated in the part of the wing where they need to be – near the front or leading or costal edge, where the pressure is the greatest. A very nice demonstration of functional morphology – adaptation of shape to function. The gray arrow

miles (560 km) N of locations in British Columbia where reported north to 56° N (Cannings 2002).

In the males identified as *L. forcipatus*, abdominal segment 2 was 3/5 as long as abdominal segment 3 (instead of 1/2 as in *L. disjunctus*), the penis vesicle was often, but not always widest near the mouth (instead of widest near the middle as in *L. disjunctus*), and the distal teeth of the cerci were much smaller than the basal teeth (instead of almost as large as the basal teeth as in *L. disjunctus*). The length of the anterior lamina viewed laterally was 1.3 - 1.5 mm in 7 males referred to *L. forcipatus* and 1.0 - 1.2 mm in several males referred to *L. disjunctus* (all from the location cited above). These latter measurements were less than reported for Ontario populations of these two species (Catling 2002) but there was nevertheless a clear separation. The males of *L. forcipatus* were more conspicuously green metallic on the dorsal abdominal segments and either green or mauve metallic on the upper sides of the thorax. One female collected had an ovipositor extending far beyond the paraprocts (further than in *L. dryas* females) and a characteristic dark spot above the metapleural carina. Although much darker than typical *L. dryas* females, this female did have metallic green on the dorsal abdominal segments.

Enallagma ebrium (Hagen), MARSH BLUET - Ten males were collected (CNC) and many others captured and released around the edges of a large pool below the escarpment on Foxholes Road west of Fort Smith (60.0517° N, -112.4783, 24° W, 24 July 2003, P. M. Catling and D. Johnson). This location is 500 miles (800 km) to the north of published records in Manitoba, Saskatchewan, Alberta and British Columbia (Walker 1953, Cannings and Stewart 1977), and apparently the northernmost record. Abdominal segments 3 - 7 were much darker in colour than in specimens from eastern Ontario.

Aeshna eremita Consuming Biting Insects

There has been some controversy over the importance of dragonflies as predators of biting insects. Habitat, location and the species of dragonflies present probably all contribute to the extent of use of biting insects as prey. More observations and quantitative studies are needed. In this connection the following observations (from field notes) may be of interest. "While I (PMC) was slowly traversing an open bog near Yellowknife on 19 July 2003, 15 individuals of *Aeshna eremita* flew in to glean mosquitoes, deer

flies, horse flies and some blackflies mostly from the top of my head but also from shoulders, back and chest. The onslaught of the biting insects appeared to lessen in intensity when the dragonflies moved in. Over a period of 10 minutes one or two dragonflies were present at least every 60 seconds. If all had not been captured, it is possible that individuals would have stayed longer and possibly consumed a large number of biting insects. In other habitats around Yellowknife where *Aeshna eremita* was present, there was no similar behaviour, so the slow movement through the open bog, and perhaps also prior experience with large mammals which also attract biting insects, may have caused the dragonflies to be less wary." More than any other dragonfly around Yellowknife, *Aeshna eremita* feeds often on deer flies and horseflies (locally collectively called "bulldogs"), but in most cases / places they capture them near people in the air rather than actually capturing those that are sitting. The foraging behaviour of dragonflies in relation to other mammals is equally interesting. Dragonflies may be significant in reducing the impact of biting insects on moose and caribou, but study will be required to determine how significant.

Aeshna eremita Adapting Foraging Behaviour to Specific Prey

It appears that *Aeshna eremita* can learn certain kinds of behaviour that is optimally effective in capturing specific kinds of prey, as evidenced by the following quote from field notes: "At a pond near Reid Lake, 62.5032° N, -113.4026° W on 27 July, Lake Darners (*Aeshna eremita*) were very numerous, with at least 5 in view at any one time. However, their flight patterns were unusual. They flew only along the shoreline, among emergent plants including *Potentilla palustris* and they were seen on 7 occasions within 10 minutes to successfully capture and eat specimens of the damselfly, *Lestes disjunctus*."

Aeshna eremita Capturing and Feeding on *Aeshna juncea*

On 27 July 2003, near Cameron River on hwy 4 at 62.5378° N, -113.8779° W, a female *Aeshna eremita* was seen to capture a male *A. juncea* in the air as the latter lifted off some low pond edge vegetation. The captor settled on a nearby shrub and proceeded to eat the almost equally sized male dragonfly first at the front of the thorax, disarticulating its head. This occurred at 7:30 on a sunny, mild evening.

Hilltopping of *Lestes congener*

On the Ingraham trail NE of Yellowknife at 62.5037° N, -114.2764° W, on 28 July 2004, 8 individuals of *Lestes congener* were found in open Aspen woods on the top of a hill and 4 were found in a similar situation on top of a nearby hill. None were found around the ponds, pools or marshes at least 200 m below the hilltops despite a search over a period of 2 hours. Hilltopping may improve chances of locating a mate and may also avoid early frost in lower lying areas. The latter may be important for relatively late flying species.

Birds Feeding On Dragonflies Hit By Cars

Along parts of hwy 3 from Yellowknife to Rae, up to 110 dead dragonflies can be seen per mile. This mostly represents those hit by cars less than a few hours earlier because they are quickly gathered by Ravens, Mew Gulls, Bonaparte's Gulls and Herring Gulls (and probably also by mammals at night). The number of dragonflies killed along a mile of road in a day probably far exceeds 100 since many are knocked clear of the pavement and the shoulder and of course many are quickly gathered by the birds. Some may also be blown off the road by wind or passing vehicles. It seems not unreasonable to speculate that a few hundred are killed along every mile of road on any single sunny day in the latter half of July at least, and perhaps throughout the summer. Of 110 specimens examined in a mile of highway on 26 July 2003, *Aeshna eremita* represented 60%, *Aeshna juncea* represented 30% and *Aeshna interrupta lineata* represented 10%. Approximately the same proportions were killed on gravel roads but the numbers were much lower, - up to 12 per mile.

Directional Movement of *Aeshna* Species

West of Yellowknife 95 % of the large dragonflies (species of *Aeshna* - see above) crossing the road in late July, crossed from the north to the south. Dispersal to the south is probably more advantageous than dispersal to the north where tundra or at least increasingly cooler climate would be encountered. On average a dragonfly, always a species of *Aeshna* and most often *Aeshna eremita*, crossed one hundred m. of road every minute. When these observations were made there was a light wind from the south.

Dependence of Endangered Bird on Dragonflies

The important role that dragonflies play in northern ecosystems is not well documented, but it has been recently found that at certain times dragonflies make up to 80 % of the food of endangered Whooping Cranes (*Grus americanus*), which breed in sloughs west of Fort Smith (Bergeson 2000). The parents gather larval forms of dragonflies, damselflies and mayflies to feed their young. (<http://www.se.gov.sk.ca/ecosystem/speciesatrisk/whoopingcrane.htm>)

Interestingly the first wild Whooping Crane to fledge in the United States (Florida) in over 60 years fed on adult dragonflies.

<http://www.pwrc.usgs.gov/whoopers/report18.htm>

Dragonfly Outing a Success

On 19 July 2003, the first NWT dragonfly outing was held at the Northern Frontier Visitors Centre on the north side of Yellowknife. It was advertised twice in the local paper and almost 100 people arrived on a warm sunny morning. A short introduction to dragonflies and to their importance was provided in the classroom, but this was kept very brief because there was not enough space for everyone to get inside the room. Next the group spent 1 1/2 hours walking around the pond on the west side of the visitor centre. The participants were a diverse group, but local families with children were predominant. There were also many retired folk and several young couples on vacation. Through the Department of Resources, Wildlife and Economic Development, nets were made available most of the approximately 25 children. A prize was offered for the one capturing the most different species. This increased interest, which was already very high. The dragonflies were captured and handled with care, identified and then released. The children brought everyone together, asking adults for help with contagious enthusiasm. Following the outing the group returned to the classroom for a question and answer wrap up. Everyone learned a lot about dragonflies and protection of environment, but more important was the experience. People, to some extent through the children, demonstrated a remarkable enthusiasm for nature and natural surroundings and a remarkable ability to work together. It was a reminder of the value of biodiversity to human health and a clear indication of the fact that people can cooperate to protect and enjoy biodiversity.

The list of species encountered on the outing was an impressive 13 for a period of 1 1/2 hours in an area of 5 acres. The species present included:

Aeshna eremita (31), *Aeshna interrupta lineata* (2), *Aeshna sitchensis* (1), *Coenagrion resolutum* (20), *Enallagma boreale* (15), *Enallagma cyathigerum* (6), *Lestes disjunctus* (200 emerging), *Leucorrhinia borealis* (8), *Leucorrhinia proxima* (6), *Libellula quadrimaculata* (30), *Somatochlora albicincta* (1), *Sympetrum obtrusum* (2 emerging), *Sympetrum internum* (1 emerging),

Dragonflies and Biodiversity Protection

A male of *Leucorrhinia hudsonica* is the central feature on the very impressive cover design of the Northwest Territories Biodiversity Action Plan. As a result of their diverse ecological relationships, vital ecosystem functions, and ease of identification, dragonflies are a particularly useful group of environmental indicators. Assessments of the general status of the 209 dragonfly species in Canada, have recently been completed to help provide a general picture of the status of Canadian Biodiversity. The results and interpretation of these assessments are expected to be available soon on the Wild Species Web site.
(<http://www.wildspecies.ca>)

Checklist of the Dragonflies of the Northwest Territories

The most recent list for the Northwest Territories (Catling 2003) included 35 species and was based on 1040 records from museum collections. This list is updated to include the additions noted above. Now 40 species are recorded for this region of Canada north of 60°. ZYGOPTERA – DAMSELFLIES

Lestidae, Spreadwings

Lestes congener Hagen, SPOTTED SPREADWING

Lestes disjunctus Selys disjunctus, COMMON SPREADWING

Lestes dryas Kirby, EMERALD SPREADWING

Lestes forcipatus Rambur, SWEETFLAG SPREADWING

Coenagrionidae, Pond Damsels

Coenagrion angulatum Walker, PRAIRIE BLUET

Coenagrion interrogatum (Selys), SUBARCTIC BLUET

Coenagrion resolutum (Selys), TAIGA BLUET

Enallagma boreale Selys, BOREAL BLUET

Enallagma cyathigerum (Charpentier), NORTHERN BLUET

Enallagma ebrium (Hagen), MARSH BLUET

Nehalennia irene (Hagen), SEDGE SPRITE

ANISOPTERA, DRAGONFLIES

Aeshnidae, Darners

Aeshna canadensis Walker, CANADA DARNER

Aeshna eremita Scudder, LAKE DARNER

Aeshna interrupta lineata Walker, VARIABLE DARNER

Aeshna juncea americana Bartenef, RUSH DARNER

Aeshna septentrionalis Burmeister, AZURE DARNER

Aeshna sitchensis Hagen, ZIGZAG DARNER

Aeshna subarctica Walker, MUSKEG DARNER

Aeshna umbrosa umbrosa Walker, SHADOW DARNER

Gomphidae Clubtails

Ophiogomphus colubrinus Selys, BOREAL SNAKETAIL

Stylurus notatus Rambur, ELUSIVE CLUBTAIL

Corduliidae Emeralds

Cordulia shurtleffi Scudder, AMERICAN EMERALD

Somatochlora albicincta (Burmeister), RINGED EMERALD

Somatochlora forcipata (Scudder), FORCIPATE EMERALD

Somatochlora franklini (Selys), DELICATE EMERALD

Somatochlora hudsonica (Hagen in Selys), HUDSONIAN EMERALD

Somatochlora kennedyi Walker, KENNEDY'S EMERALD

Somatochlora minor Calvert in Harvey, OCELLATED EMERALD

Somatochlora sahlbergi Trybom, TREELINE EMERALD

Somatochlora septentrionalis (Hagen), MUSKEG EMERALD

Libellulidae, Skimmers

Leucorrhinia borealis Hagen, BOREAL WHITEFACE

Leucorrhinia hudsonica (Selys), HUDSONIAN WHITEFACE

Leucorrhinia patricia Walker, CANADA WHITEFACE

Leucorrhinia proxima Calvert, VARIABLE WHITEFACE

Libellula quadrimaculata Linnaeus, FOUR-SPOTTED SKIMMER

Sympetrum costiferum (Hagen), SAFFRON-BORDERED MEADOWFLY

Sympetrum danae (Sulzer), BLACK MEADOWFLY

Sympetrum internum Montgomery, CHERRY-FACED MEADOWFLY

Sympetrum obtrusum (Hagen), WHITE-FACED MEADOWFLY

Sympetrum madidum Hagen, RED-VEINED MEADOWHAWK

ACKNOWLEDGEMENTS

Anne Gunn kindly provided specimens collected from the Yellowknife area and from McKay Lake area and also assisted in field studies near Yellowknife. Doug Tate supplied specimens of 8 species from Rabbitkettle Lake in Nahanni National Park Reserve. Richard Popko supplied specimens of three species of *Aeshna* from Norman Wells. Some Coenagrionid larvae from the Nahanni area supplied by Dwayne Lepitzki appeared to have characteristics of *Ischnura* but could not be conclusively identified. The closest record of an *Ischnura* to NWT is that of *I. damula* at Liard River in northwestern British Columbia.

LITERATURE CITED

Bergeson, D. G., M. Bradley, and G. Holroyd. 2000. Food items and feeding rates for wild Whooping Crane colts in Wood Buffalo National Park. Pp. 36-39 in D. H. Ellis. Ed., Proceedings of the eighth North American Crane workshop.

Cannings, R. A. 2002. Introducing the dragonflies of British Columbia and the Yukon. Royal British Columbia Museum. 675 Belleville St. Victoria BC V8W 9W2. 96 pp.

Cannings, R. A. And K. M. Stuart. 1977. Dragonflies of British Columbia. British Columbia Provincial Museum, Handbook No. 35. Victoria. 254 pp.

Cannings, S.G. and R.A. Cannings. 1997. Dragonflies (Odonata) of the Yukon. Pp. 169-200 in H.V. Danks and J.A. Downes (eds.), Insects of the Yukon. Biological Survey of Canada (Terrestrial Arthropods), Ottawa. 1034 pp.

Catling, P. M. 2002. Evaluation of some characters separating male *Lestes disjunctus* and *Lestes forcipatus* in Ontario. Pp. 51-58 in P. M. Catling, C. D. Jones and P. Pratt, eds., Ontario Odonata, vol. 3. Toronto Entomologists Association, Toronto, Canada.

Catling, P. M. 2003. Dragonflies (Odonata) of the northwest territories - status ranking and

preliminary atlas. Department of Resources, Wildlife and Economic Development Government of the Northwest Territories. 49 pp.

Hutchings, G. E. 2002. A list of the Odonata of Athabasca Sand Dunes Provincial Wilderness Park, Saskatchewan. Argia (The News Journal of the Dragonfly Society of the Americas) 14(4): 5-8.

Needham, J.G., M.J. Westfall, Jr., and M.L. May. 2000. Dragonflies of North America. Scientific Publishers, Gainesville, Florida. 940 pp.

Walker, E.M. 1953. The Odonata of Canada and Alaska, volume 1, part 1: General. part 2: The Zygoptera - Damselflies. University of Toronto Press. 292 pp.

Walker, E.M. 1958. The Odonata of Canada and Alaska, volume 2, part III: The Anisoptera - four families. University of Toronto Press. 318 pp.

Walker, E.M. and P.S. Corbet. 1975. The Odonata of Canada and Alaska. Volume 3: The Anisoptera - Three families. University of Toronto Press, Toronto. 308 pp.

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

LAWSUIT FILED TO ESTABLISH HABITATS FOR DRAGONFLY

Dan Egan <degan@journalsentinel.com>
From the Feb. 6, 2004 editions of the Milwaukee Journal Sentinel. Posted: Feb. 5, 2004

A coalition of conservation groups has filed suit in federal district court in Washington, D.C., over its claims the federal government is not doing enough to help the endangered Hine's emerald dragonfly.

The green-eyed bug can be found in pockets of eastern Wisconsin as well as Michigan, Illinois and Missouri. It has disappeared from its historic range in Ohio, Indiana and Alabama because of habitat loss, according to the conservationists filing suit.

The lawsuit, filed Wednesday, seeks to force the U.S. Fish and Wildlife Service to formally designate areas as "critical habitat" for the beleaguered species, something that hasn't happened despite its 1995 listing on the federal Endangered Species Act.

"We don't want this species sitting on the Endangered Species Act in perpetuity, and when the Fish and Wildlife Service fails to designate

After breakfast, we checked out two pools along US 1 in the town of Big Pine Key. Almost immediately, we saw the red *Orthemis* patrolling around the edges of two borrow pits. We both got several males and females. The males are identical to the single red *Orthemis* male I got in the Dominican Republic. I also got a lone purple *Orthemis ferruginea*, which is similar to the purple *Orthemis* I got in the Dominican Republic. Ken reared a male of the red *Orthemis* and a female of *Tramea onusta* from one of the small ponds.

We then drove up the sunny, scenic Keys to get to the Everglades to do some collecting before nightfall. At a canal along Hwy. 997, we got teneral *Nehalennia pallidula*, *Epitheca stella*, *Erythrodiplax umbrata*, and a purple *Orthemis ferruginea*. That evening, we stayed at the Everglades City motel again and had conch fritters and Key Lime pie at the nearby Oar House restaurant.

The next morning we collected some *Miathyria marcella* larvae hiding among the roots of the hyacinths and water lettuce in a canal. Then, we headed out for Sarasota to look for *Crocothemis servilia* larvae where I had seen the adults a couple of years before. When we got to the 7-Eleven retention pond, I could not believe my eyes! Several males and a female were flying around the pond! I hurried back and told Ken while picking up my net. We each got one, then we settled down to

The next morning, we were greeted by very cold weather in Tallahassee as we set out to collect some *Enallagma davis* larvae nearby. Ken was successful in getting a few in the deeper water of Dog Pond plus we got some *Anax longipes*, *Progomphus bellei*, and *Gomphus cavillaris brimleyi* larvae from nearby sandy lakes. The following day, we drove to Hosford to look for *Gomphaeschna antilope* larvae in Whitehead Bay and vicinity where I saw dozens of adults in early April several years earlier. No such luck! The area is extensive and there was too much habitat for the larvae to hide in. I can understand why it is such an elusive species in the larval stage! We did do a lot better at nearby Bear Creek where Ken got larvae of *Cordulegaster maculata*, *Gomphus geminatus*, *Stylurus ivae*, and *S. laurae*. Ken took off the next morning to do some collecting near Torreya State Park on his way back to Alabama.

All in all, it was a very successful trip! The warm, sunny Florida Keys was a welcome respite from the cold winter weather in northern Alabama and north Florida. As an extra bonus, we did not get any mosquito or "no-see-um" bug bites in the Everglades or on the Keys. We hope to do it again soon and continue to inventory the Odonata of the National Key Deer Refuge under their supervision. Now that we know the red *Orthemis* is a permanent resident in the Keys, hopefully we can study it and learn more about this new dragonfly to the United States, especially its taxonomic status.

FIRST RECORDS OF PACIFIC SPIKETAIL, *CORDULEGASTER DORSALIS*, IN IDAHO

Kent Fothergill, John Keebaugh, and Miriam Austin

Red Willow Research, Inc. 780 Falls Avenue #390, Twin Falls, ID 83301 (KF & MA); Orma J. Smith Museum of Natural History, Albertson College of Idaho, Caldwell, ID 83605 (JK); <cyberzono@netscape.net> (KF)

Donnelly (2004) has compiled a county-based atlas of North American Odonata and reports no occurrence of *Cordulegaster dorsalis* in Idaho. Kondratieff (2000) compiled a United States Odonate atlas based on verified specimens, and also reports no occurrences of *C. dorsalis* in Idaho. The Slater Museum of Natural History (Paulson 1997) compiled a list of Idaho Odonata based on occurrences in literature (with much of the Idaho information coming from Logan (1967)) and field sampling, and again reports no occurrence of *C. dorsalis* in Idaho. Finally the Idaho Museum of Natural History's (2002) Digital Atlas of Idaho also has no occurrence of *C. dorsalis* in Idaho.

Needham, Westfall and May (2000) list *C. dorsalis* as occurring in the Western Great Basin and Pacific Northwest both biotic provinces with which Idaho has strong affinities. Based on this information Dunkle (2000) includes portions of Idaho in his range map for *C. dorsalis* (Dunkle 2004).

C. dorsalis has been netted in Ada and Cassia counties. The Ada county occurrence was on June 29, 2003 during a North American Butterfly Association Fourth of July butterfly count above Dry Creek near Bogus Basin road and was viewed by many members of the Idaho Entomology Group (IEG). *C. dorsalis* adults were observed cruising between scattered hackberries (*Celtis*). A specimen was kept, but is no longer in existence.

The Cassia county netted occurrence was part of an Odonate survey performed July 22, 2003 (RWR 2003). A female was observed ovipositing in Rock Creek at Harrington Fork and a male was netted cruising within the dense dogwood/water birch (*Cornus/Betula*) riparian zone. The netted insect was identified and released. Results were of the survey were sent as a courtesy to Dennis Paulson who inquired as to whether a specimen was obtained and informed the authors of the unique nature of these sightings. On July 19, 2003 *C. dorsalis* adults were patrolling the pinyon/juniper (*Pinus/Juniperus*) forest edges near the Jim Sage Canyon spring enclosure area in Cassia county. Digital photographs of *C. dorsalis* adults taken at this time were e-mailed to Dennis Paulson who

identified the photographs as *Cordulegaster dorsalis dorsalis* (Paulson 2003).

A sight record of adults exists for Elmore county, May 24, 2002 at the intersection of Bennet Mountain Road and Little Canyon Creek; (Keebaugh 2002) which is the first reported occurrence in Idaho. The habitat utilized here was a willow/ chokecherry (*Salix/Prunus*) riparian zone.

C. dorsalis is now also documented by photographic evidence in Cassia county, and by sight in Ada and Elmore counties. Given the wide range of habitats and the long flight period (May 24 to July 22) in which this charismatic dragonfly was found, it seems quite likely that *C. dorsalis* will be found in other locations in Idaho.

REFERENCES

- Donnelly, T. W., 2004, Distribution of North American Odonata. Part I: Aeshnidae, Petaluridae, Gomphidae, Cordulegastridae, Bulletin of American Odonatology: 7(4):61-90
- Dunkle, S. W., 2000, Dragonflies through Binoculars, Oxford University Press, New York NY, 266pp
- Dunkle, S. W., 2004, Personal Communication
- Idaho Museum of Natural History, 2002, Digital Atlas of Idaho <http://www.imnh.isu.edu/digitalatlas>
- Logan, E.R., 1967, The Odonata of Idaho, an unpublished M.S. thesis, University of Idaho, Moscow ID,
- Keebaugh, J., 2002, Field Notebook
- Kondratieff, B. C. (coordinator), 2000, Dragonflies and Damselflies (Odonata) of the United States. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page. <http://www.npwrc.usgs.gov/resource/distr/insects/dfly/dflyusa.htm> Version 12DEC2003).
- Needham, J. G., M.J. Westfall Jr., M.L. May., 2000, Dragonflies of North America, Scientific Publishers, Gainesville FL, 939pp
- Paulson, D. R., 1997, The Odonata of Idaho, Slater Museum of Natural History, University of Puget Sound, Tacoma, WA <http://www.ups.edu/biology/museum/IdahoOD.htm>

Paulson, D. R., 2003 Personal Communication

Red Willow Research, Inc., 2003, Magic Mountain
Dragonfly Count,
<http://www.redwillowresearch.com>

== == == == == == == == == == == == == == == ==

NEW COUNTY RECORDS FOR IOWA - 1999 TO 2003

Steve Hummel, P. O. Box 121, Lake View, IA
51450

Since the publication of the Odonata of Iowa by Robert W. Cruden and O. J. Gode (2000), fairly intensive collecting has continued in Iowa. Most of the additions included here are records from the 1999 to 2003 collecting seasons, with a few older records.

Two species, *Somatochlora linearis* and *Libellula incesta*, were reported as new to the state by Ann Johnson (2003). However, Johnson did not include information on the counties the records were from. Both species were discovered in July 2002.

S. linearis was originally listed for Iowa based on a possibly misidentified specimen. Loudon (1933) reported *S. linearis* from the collection at Iowa State University for Story Co. *S. linearis* for Iowa was included in Needham, Westfall, and May (2000), but Cruden and Gode (2000) listed it as excluded or questionable on the basis that a specimen in the Iowa State University collection had also been identified as *S. ensigera* and *S. tenebrosa*. In August 1978, in Tama Co., Hal White (unpublished data) recorded a large *Somatochlora* ("large as *cingulata*. Perhaps *ensigera* or *linearis* two good but unsuccessful swings. Flying with *A. junius*"). Aaron Brees found this species in Wayne Co in July of 2002. In August, also 2002, I got a good sight record in Madison Co. as well. This represents a northern and western extension in the known range of this species based on the range map in Dunkle (2000).

L. incesta has not been listed previously from the state. Again, Aaron Brees first found this species, also in Wayne Co., in July or 2002. Based on the distribution for the species given in Needham, Westfall, and May (2000) and the distribution map in Dunkle (2000), this Iowa record fills in a gap for the species.

A third species, *Sympetrum danae*, had also been removed from the state list by Cruden and Gode (2000). The only source of this species presence in Iowa was Montgomery's (1967) checklist. Cruden and Gode were unable to discover a locality record for the species. I also had been unable to locate any record for the species. However, a specimen was captured in Sac Co. in September, 2001.

Gomphus ventricosus is another species Cruden and Gode (1998) excluded from the state list. They questioned the inclusion of the species by Hummel and Haman (1975, 1977). Our inclusion of the species was based on a female specimen collected by Miller on 6/7/1906 in Black Hawk Co., and identified by Kormondy (1960) from a female specimen in the Williamson-Kennedy Odonata collection of the University of Michigan Museum of Zoology. Cruden and Gode (1998) correctly stated that Miller (1906) did not list the species (he did list "*Gomphus* sp.? Two females. These are teneral, and Mr. Williamson hesitates to give them a specific name."), but, did not have the Kormondy reference. Beckemeyer (2002) lists *G. ventricosus* for Iowa, Minnesota, and Missouri. While it has not been collected in Iowa since 1906, it could be found along some of the larger rivers particularly in eastern and northeastern Iowa.

Arigomphus cornutus should be removed from Muscatine Co. Cruden and Gode (2000) misread a data point on a map in Hummel and Haman (1977). The data point was for *Gomphus crassus* (Wells, 1917, who referenced Wilson and gave a date of 1916, but Wilson (1920) does not list the species from Fairport). *A. cornutus* has not yet been collected in Muscatine Co. Cruden and Gode (2000) excluded *G. crassus* from the Iowa list.

The combination of earlier workers (Elrod 1898, Miller 1906, Wilson 1909 and 1920, Whedon 1912, Wells 1917, Hoffman 1924, Yeager 1932, Loudon 1933, Hummel and Haman 1975 and 1977, and Hummel 1978), Cruden and Gode, and the information presented here, has produced an average number of 47 (46.5758) species per county in Iowa. The five counties with the greatest number of species per county are Black Hawk, Johnson, Muscatine, and Story with 69 each, and Sac with 67. The five counties with the fewest number of species per county are Mitchell with 31, Adams, Humboldt, and Fremont with 33 each, and Montgomery with 34. The counties with the greatest numbers of species, not surprisingly, are, or were, the residence counties of Robert W.

Cruden and O. J. Gode (Johnson) or Steve Hummel (Black Hawk and Sac).

Western, southern, and northeastern Iowa are especially in need of additional collecting as these areas have been under collected in the past and they have a potential of additional species for the state. Western Iowa could harbor species typically found farther west (Beckemeyer, 2002), and northeast Iowa could have several species reported from Wisconsin (Legler, Legler, and Westover, 1998). Southern Iowa has been the source of 2 of the 3 new state records reported here. Additional species known from northern Missouri (Trial and Belshe, 2002) could also be found in southern Iowa.

In 2001 Ann Johnson set up a website using the data from Cruden and Gode (2000) and included distribution maps and county species lists. As new county records have been discovered, these species have been added to the maps and species.

This update includes 241 new county records for 71 species in 53 of Iowa's 99 counties and is based on the work of Steve Hummel (SH), Ann Johnson (AJ), Bob Cruden (BC), and Aaron Brees (AB), plus others as named in the list. Some records are also from specimens that were identified by the author from the Iowa State University (ISU), Iowa Wesleyan College (IWC), and Iowa Lakeside Laboratory (IaLL) collections.

(species, county, collector, date); (NSR) = new state record

Calopteryx aequabilis: Carroll, SH, 8/3/1999; Crawford, BC, 7/5/2001

Hetaerina titia: Appanoose, AB, 8/26/2002; Audubon, AB & AJ, 8/24/2003; Delaware, BC, 10/3/2001; Dubuque, BC, 10/3/2001; Story, AJ, 9/3/2001; Wayne, AB, 8/17/2002

Archilestes grandis: Boone, W. S. Craig, 9/17/1962 (ISU); Ida, SH, 8/12/2003; Marion, AB, 9/16/2002

Lestes disjunctus australis: Boone, D. L. Moeller (ISU), 5/3/1965; Carroll, SH, 6/17/2001; Greene, SH, 5/19/2001; Ida, SH, 6/15/2001; Lucas, AB & AJ, 7/6/2002; Warren, AJ, 6/15/2003; Wayne, AB & AJ, 7/6/2002;

Lestes dryas: Cerro Gordo, AJ, 6/30/2001

Lestes eurinus: Cedar, BC, 6/11/2001; Fayette, AB & AJ, 7/21/2003

Lestes forcipatus: Cedar, BC, 6/11/2001

Lestes inequalis: Fayette, AB & AJ, 7/21/2003

Lestes rectangularis: Carroll, SH, 6/30/1999; Greene, SH, 7/16/2001; Howard, SH, 8/1/2002; Ida, SH, 6/15/2001; Winneshiek, SH, 8/2/2002

Lestes unguiculatus: Webster, SH, 7/9/2002

Amphiagrion sp: Lucas, AB, 9/4/2002; Palo Alto, SH, 6/21/2001; Polk, J. Laffoon (ISU), 5/26/1947; Story, L. S. Wells (ISU), 6/13/1916; Warren, AJ, 7/1/2001

Argia alberta: Shelby, BC, 7/5/2001

Argia apicalis: Palo Alto, SH, 7/11/2003; Shelby, BC, 7/7/2001; Wright, SH, 8/14/2003

Argia fumipennis violacea: Howard, SH, 8/3/2002; Shelby, BC, 7/5/2001

Argia moesta: Appanoose, AB & AJ, 7/4/2003; Audubon, AB & AJ, 8/24/2003; Boone, M. A. Zwanzinger (ISU), 9/25/1963; Cerro Gordo, AJ, 6/30/2001; Crawford, SH, 8/19/2001; Harrison, AJ, 7/7/2001; Wayne, AB, 7/20/2002

Coenagrion angulatum: Emmett, H. Weston (ISU), 5/25/1949; Story, ISU, 5/20/1929

Coenagrion resolutum: Hancock, H. Borchers (ISU), 6/24/1966; Story, L. S. Wells (ISU), 7/3/1916

Enallagma anna: Floyd, BC, 6/26/2001; Howard, SH, 8/1/2002

Enallagma antennatum: Ida, SH, 8/25/2003

Enallagma aspersum: Henry, Shriner (IWC), 7/16/1947; Ida, SH, 6/15/2001; Wayne, AB & AJ, 7/6/2002; Webster, SH, 7/9/2002

Enallagma basidens: Buena Vista, SH, 8/19/2002

Enallagma carunculatum: Mahaska, AJ, 7/4/2001

Enallagma cyathigerum: Carroll, SH, 7/21/1999

Enallagma ebrium: Henry, IWC, 10/5/1971; Story, L. S. Wells (ISU), 7/6/1916

Enallagma geminatum: Cerro Gordo, AJ, 9/7/2002; Ida, SH, 5/29/2003

Enallagma signatum: Buena Vista, SH, 6/11/2000; Carroll, SH, 6/30/1999; Ida, SH, 8/14/1999; Polk, AJ, 5/29/2001; Webster, SH, 7/7/2001

Enallagma traviatum: Pottawattamie, AB & AJ, 7/19/2003; Warren, AJ, 7/1/2001

Enallagma vesperum: Calhoun, SH, 7/22/2002; Madison, Roslea I. Johnson, 9/12/2003.

Ischnura hastata: Carroll, SH, 6/8/1999

Ischnura posita: Sac, SH, 6/3/2002; Story, ISU, 6/5/1927

Nehalennia irene: Allamakee, SH, 7/15/2000; Boone, D. R. Miller (ISU), 7/2/1963; Carroll, SH, 6/30/1999; Monona, S. M. Gaud (ISU), 6/6/1960; Webster, SH, 7/9/2002

Aeshna constricta: Calhoun, SH, 9/16/2000; Marion, AJ, 9/6/2003; Worth, AB & AJ, 9/7/2002; Wright, AB & AJ, 9/2/2002

Aeshna multicolor: Allamakee, SH, 7/17/2003; Dallas, AJ, 8/2/2003; Fayette, AB & AJ, 7/21/2003; Monona, AB & AJ, 6/29/2003

Aeshna umbrosa: Carroll, AJ, 9/20/03; Dubuque, BC, 10/3/2001; Ida, SH, 8/21/2003; Polk, IWC, 7/??/1931; Wayne, AB, 7/25/2002

Boyeria vinosa: Calhoun, SH, 8/29/1999; Webster, SH, 9/7/2002

Epiaeschna heros: Cedar, BC, 6/1/2002

Nasiaeschna pentacantha: Lucas, AB & AJ, 7/4/2003; Wayne, AB & AJ, 7/13/2002

Arigomphus cornutus: Clayton, BC, 6/10/2001; Dallas, AB, 6/23/2003; Sac, SH, 6/26/2001; Warren, AJ, 6/14/2003

Arigomphus submedianus: Dallas, AJ, 8/2/2003; Hamilton, SH, 6/29/2001; Harrison, AJ, 7/7/2001; Monona, AB & AJ, 6/29/2003; Polk, AB & AJ, 6/28/2003; Pottawattamie, AJ, 6/30/2002; Shelby, BC, 7/5/2001

Dromogomphus spoliatus: Dallas, AJ, 8/2/2003; Linn, AJ, 8/1/2001; Wayne, AB, 7/13/2002

Gomphus externus: Audubon, AB & AJ, 8/24/2003; Franklin, BC, 7/2/2002; Madison, AJ, 6/16/2002; Monona, AB & AJ, 6/29/2003; Pottawattamie, AB & AJ, 7/19/2003; Shelby, BC, 7/5/2001; Warren, AJ, 5/19/2002

Gomphus graslinellus: Wayne, AB, 7/20/2002

Gomphus militaris: Decatur, AB & AJ, 7/3/2003; Monona, AB & AJ, 6/29/2003

Gomphus vastus: Calhoun, SH, 6/23/2001; Dallas, AB, 6/23/2003; Hamilton, SH, 8/27/2003; Mills, Loren Padelford, 6/9/1999; Story, Erwin Klaas, 7/13/2003; Webster, SH, 7/7/2001

Ophiogomphus rupensulensis: Winneshiek, SH, 7/17/2003

Progomphus obscurus: Audubon, AJ, 8/24/2003; Buchanan, BC, 7/2/2002; Calhoun, SH, 7/8/2002; Carroll, SH, 8/3/1999; Crawford, BC, 7/5/2001; Hamilton, SH, 8/27/2003; O'Brien, SH, 7/6/2002; Webster, SH, 10/13/2001; Wright, SH, 8/27/2003

Stylurus amnicola: Boone, L. S. Wells (ISU), 7/14/1916; Crawford, SH, 8/19/2001; Greene, SH, 7/16/2001; Ida, SH, 7/28/2001; Wright, SH, 8/27/2003

Stylurus notatus: Calhoun, SH, 7/8/2002; Cerro Gordo, AB & AJ, 9/7/2002; Wright, SH, 8/27/2003

Stylurus plagiatus: Appanoose, AB, 8/26/2002; Dubuque, IL St. Mus. Coll.; Greene, SH, 7/16/2001; Wayne, AB, 7/20/2002

Macromia illinoensis: Appanoose, AB, 8/26/2002; Audubon, AB & AJ, 8/24/2003; Bremer, AJ, 6/23/2002; Dallas, AJ, 8/11/2003; Marion, AB, 9/13/2002; Story, SH 7/24/2003; Warren, AJ, 6/15/2003; Wayne, AB & AJ, 7/6/2002; Wright, SH, 8/14/2003

Epithea cynosura: Carroll, SH, 6/8/1999; Lyon, IaLL, 6/13/1973

Somatochlora ensigera: Wright, AJ, 9/2/2002

Somatochlora linearis (NSR): Madison, SH, 8/24/2002; Wayne, AB & AJ, 7/13/2002

Celithemis elisa: Ida, SH, 7/28/2001; Lucas, AB, 8/14/2002; Story, Stephanie J. Black, 7/18/2002; Wayne, AJ, 6/21/2003

- Celithemis eponina*: Buena Vista, SH, 7/14/2001; Carroll, SH, 7/31/1999; Decatur, SH, 7/15/2002; Ida, SH, 7/28/2001; O'Brien, SH, 7/6/2002; Shelby, BC, 7/5/2001
- Leucorrhinia intacta*: Ida, SH, 7/11/2003; Warren, AJ, 5/18/2002
- Libellula incesta* (NSR): Wayne, AB, 7/20/2002
- Pachydiplax longipennis*: Crawford, BC, 7/5/2001; Ida, SH, 7/29/2001
- Pantala flavescens*: Appanoose, AB & AJ, 7/4/2003; Buena Vista, SH, 7/11/2003; Ida, SH, 8/14/1999; Wayne, AB & AJ, 7/6/2002; Winneshiek, SH, 8/2/2002; Worth, AB & AJ, 9/7/2002
- Pantala hymenaea*: Dallas, AB & AJ, 6/28/2003; Decatur, AB & AJ, 7/3/2003; Ida, SH, 8/14/1999; Marion, AB, 9/20/2002; Monona, AB & AJ, 6/29/2003; Story, Stephanie J. Black, 8/7/2002; Wayne, AB, 7/21/2002
- Sympetrum ambiguum*: Lucas, AB & AJ, 8/31/2002; Wayne, AB, 7/20/2002
- Sympetrum corruptum*: Boone, AJ, 6/16/2002; Franklin, AB & AJ, 7/12/2003; Greene, SH, 6/22/2000; Harrison, AJ, 7/7/2001; Henry, Eastman (IWC), 4/14/1937; Howard, SH, 8/3/2002; Wayne, AB, 4/15/2003
- Sympetrum costiferum*: Marion, AB, 10/7/2002; Webster, SH, 7/9/2002; Wright, AJ, 9/19/2003
- Sympetrum danae*: (NSR) Sac, SH, 9/26/2001
- Sympetrum internum*: Calhoun, SH, 9/16/2000; Carroll, SH, 6/17/2001; Webster, SH, 10/5/2002; Winnebago, IWC, 6/21/1932
- Sympetrum obtrusum*: Calhoun, SH, 6/23/2001; Webster, SH, 10/5/2002
- Sympetrum occidentale fasciatum*: Buena Vista, SH, 8/19/2002; Howard, SH, 8/3/2002; Lucas, AJ, 9/8/2002; Mahaska, AJ, 7/4/2001; Story, Jennifer Smith, 7/15/2001; Worth, AJ, 9/7/2002
- Sympetrum rubicundulum*: Palo Alto, SH, 7/11/2003
- Sympetrum vicinum*: Carroll, SH, 8/3/1999; Clay, SH, 8/19/2002; Humboldt, SH, 8/14/2003; Lucas, AB, 9/4/2002; Story, Stephanie J. Black, 9/5/2002; Webster, SH, 10/5/2002
- Tramea lacerata*: Calhoun, SH, 6/23/2001
- Tramea onusta*: Cerro Gordo, AJ, 6/30/2001; Crawford, BC, 7/5/2001; Dallas, AJ, 6/28/2003; Decatur, AB & AJ, 7/3/2003; Des Moines, Atwood (IWC), 6/3/1931; Greene, SH, 6/22/2000; Harrison, AJ, 7/7/2001; Ida, SH, 6/15/2001; Lucas, AB, 9/4/2002; Monona, AB & AJ, 6/29/2003; Pottawattamie, AB & AJ, 7/19/2003; Story, AJ, 7/28/2002; Wayne, AB & AJ, 7/6/2002; Webster, SH, 7/9/2002
- I would like to thank Ann Johnson for access to the database of her web site (<http://www.iowaodes.com>) as a source for many of the county records, especially those of Ann Johnson, Aaron Brees, and the others named in the list. In addition, Ann Johnson and Aaron Brees were kind enough to check on the accuracy of their records as reported here.
- Literature cited:
- Beckemeyer, Roy J. 2002. Odonata In The Great Plains States: Patterns of Distribution and Diversity. Bull. of Am. Odonatology 6(3):49-99.
- Cruden, Robert W., and O. J. Gode. 1998. Iowa's Odonata, Declining and/or Changing? Proc. Iowa Acad. Sci. 105(2):67-81.
- Cruden, Robert W., and O. J. Gode. 2000. The Odonata of Iowa. Bull. of Am. Odonatology 6 (2):13-48.
- Dunkle, Sidney W. 2002. Dragonflies Through Binoculars. Oxford University Press. 266 pp.
- Elrod, M. J. 1898. Iowa Odonata. Ent. News. 9:7-10.
- Hoffman, C. 1924. Odonata (dragonflies and damselflies) of Henry County. Proc. Iowa Acad. Sci. 31:441-442.
- Hummel, M. S. 1978. A Checklist of the Odonata of Dickinson County, Iowa. Cordulia 4:127-128.
- Hummel, M. S. and A. C. Haman. 1975. Notes on the Odonata of Black Hawk County, Iowa. Ent. News (86):63-64.

collections were made over the following eight days, indicating that a small population was present. Exuviae were found on several days at this site again in 2003. In addition, a single exuvia was collected at a location on Mactaquac Lake on 25 June 2003 by A.W.T. These collections are the first reported for Canada. *N. obsoleta* has been reported from numerous localities in Maine (Brunelle 1999b), the closest being Crawford, Washington Co., approximately 125 km from the Fredericton site.

New Provincial Records

***Gomphus vastus*:** A.W.T. encountered large numbers of adults on the south shore of the Saint John River site at Fredericton on 11 August 2001. Collections of exuviae and adults by the authors during the next two years indicated that *G. vastus* was the most abundant Anisoptera species in the lower (tidal) reaches of the Saint John River and its tributaries. It is also present in Mactaquac Lake, a large headpond formed by a dam located approximately 2 km upstream from the head-of-tide and 17 km west of Fredericton, and in Indian Lake and Grand Lake, 25 and 45 km east of Fredericton. Searches of other river systems in southern New Brunswick have failed to turn up additional sites for the species. *G. vastus* seems to have a sporadic distribution in the northeast. It is known from a few rivers in Maine; the closest known site to the New Brunswick occurrences is 150 km to the southwest on the Penobscot River at Mattamiscontis Township (Brunelle 2001).

***Aeshna clepsydra*:** On 14 August 2002, Mary Sabine briefly observed an ovipositing female damselfly that she identified as *A. clepsydra* at Waltons Lake, Kings Co (45.4639(N, 66.0390(W)). Approximately 30 minutes later D.L.S collected a male several hundred metres distant. Walton Lake is a shallow, gravel-bottomed lake with a dense growth of *Juncus militaris* in the near-shore area. Subsequent to this discovery, the authors and others have collected *A. clepsydra* at a number of sites in southwestern New Brunswick. The species is known from scattered locations throughout adjacent Maine and southwestern Nova Scotia, so its discovery in New Brunswick was not surprising.

***Arigomphus furcifer*:** H.S.M. collected an adult female at Jemseg, Queens Co. (45.8243(N, 66.1173(W)) on 30 June 2003. The site consisted of an elevated gravel road that crossed an extensive floodplain area between the Saint John and Jemseg Rivers. Numerous ponds and rich marshes are

found throughout the area adjacent to the roadbed. A.W.T. collected another adult female at the site on 7 July 2003. Ample suitable habitat exists throughout the floodplain of the lower Saint John River; future search for the species will probably yield new sites. *A. furcifer* is known from scattered locations in southern Maine. The New Brunswick record extends the northeastern limits of the species range approximately 110 km east of the nearest locality reported in Brunelle 1999b.

***Hetaerina americana*:** On 8 September 2003, D.L.S. encountered 100-150 adults on the Eel River at Maxwell, York Co. (45.9284(N, 67.7038(W)). Later that day and on several other dates during the succeeding week the species was encountered by the authors and Kate Bredin (Atlantic Canada Conservation Data Centre) on the Eel River as well as at a single site on the Magaguadavic River, 40 km to the southeast. The habitat of at these sites was fast-flowing water with cobble and/or bedrock substrate. This species is known from several sites in Maine, the closest being at the Penobscot River approximately 75 km southwest of the Eel River (Brunelle and deMaynadier 2000). *H. americana* was recently rediscovered in Maine: the last collection in the state had occurred in 1901 (Brunelle 1999b). The lack of records in Maine and in New Brunswick is likely due to its late flight season, which starts well after most odonate survey activity has ceased. There are several other rivers in central and southern New Brunswick with suitable habitat and that deserve further survey for this species.

First New Brunswick or Canadian records of *G. abbreviatus* and *G. vastus* are in the collection of A.W.T. Subsequent collections of these species by the authors, as well as initial collections of the remaining four species, are in the collection of the New Brunswick Museum.

We thank Stuart Tingley for stimulating our interest in the Odonata of the province and Paul Brunelle for confirming D.L.S.'s first exuvial records of *N. obsoleta*.

REFERENCES

- Brunelle, P.M. 1997. Distribution of dragonflies and damselflies (Odonata) of the Atlantic Provinces, Canada. *Northeastern Naturalist* 4(2):61-82.
- Brunelle, P.M. 1999a. Additions to the lists of dragonflies (Odonata: Anisoptera) of the Atlantic

Provinces, Canada. Northeastern Naturalist 6(1):35-38.

Brunelle, P.M. 1999b. Distribution of damselflies and dragonflies (Odonata) of Maine, United States. Northeastern Naturalist 6(2):95-118.

Brunelle, P.M. and P. deMaynadier. 2000. The 1999 Season. Mainensis 1(1): 2-3.

Brunelle, P.M. 2001. The 2000 Season. Mainensis 2(1): 7-8.

Needham, J.G., M.J. Westfall, Jr., and M.L. May. 2000. Dragonflies of North America. Gainesville, Scientific Publishers. 939 pp.

Walker, E.M. 1953. The Odonata of Canada and Alaska, Vol. 1. Toronto, University of Toronto Press. 292 pp.

Walker, E.M. 1958. The Odonata of Canada and Alaska, Vol. 2. Toronto, University of Toronto Press. 318 pp.

Walker, E.M., and P.S. Corbet. 1975. The Odonata of Canada and Alaska, Vol. 3. Toronto, University of Toronto Press. 307 pp.

Westfall, M.J., Jr., and M.L. May. 1996. Damselflies of North America. Gainesville, Scientific Publishers. 649 pp.

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|

ANAX AMAZILI NEW FOR GUADELOUPE AND NOTES ON OTHER RARE SPECIES (FRENCH WEST INDIES)

François Meurgey and Carolyne Perron
Museum d'Histoire Naturelle de Nantes (France)
<Francois.meurgey@mairie-nantes.fr>

During our second mission to Guadeloupe (from April 21 to May 11, 2003) we returned to Vauchelet ponds to verify the *Ischnura hastata* population. Since our last visit in 2002, these ponds were turned into shrimp's rearing basins, which are now the private property of a restaurant. After a little disappointment, we found a great population of *Ischnura hastata*, and discovered a single *Anax amazili* exuvium hanged upon the chain link fence surrounding the ponds. We observed some Aeshnidae inside the surrounding fence, but did not get authorization to pass through the gate to capture them. These two 100 square meters basins are close to each other, separated by a slope, and fed by two

streams. There is no heliophytic vegetation along the banks or any submersed plants in the water. To our knowledge, there is only one record of this species in the French West Indies, in Martinique (Tsuda, 1991), where it seems that the species was not seen since then. This seems to be the first observation of *A. amazili* in the Guadeloupe archipelago and presence of an exuvium proves that this species reproduces, but perhaps not regularly.

Again with *Anax*. We observed a female *A. concolor* egg laying in submersed vegetation under mangrovs, during the survey of the Etang Zombis (29 April 2003) at 420 meters high. We collected 67 exuviae along the banks of this pond. *Anax concolor* has been known to Guadeloupe since 1992, but in small numbers, and only adults without any reproductive behaviour were observed. Exuviae were collected on young trees, between 50 cm and 3,50 meters high. Access to Etang Zombis is hazardous; mangroves surround the water and rainforest is very dense around the pond. There is no road access, which could explain why this site had never been surveyed.

Tramea binotata is another rare species in Guadeloupe archipelago. Known from Basse-Terre since 1997 (Hoffman, 1999) the observations are only of imagos (two males on the Goyave river), without additional information. Three distinct populations were observed on three different sites: Grand Etang, Etang Zombis and Vauchelet's rearing basins. We collected many exuviae along the banks of these ponds and two newly emerged *Tramea* were caught. Curiously enough, this species was never seen before 1997 on the island, even at the Grand Etang, a well known naturalists site; is this a migrant species that can regularly breed on the island? These observations prove that this species is reproductive on Guadeloupe, at least in the 2003 season. One of the 2004 mission objectives is to verify the persistence of these three populations.

References

- Hoffman, C., 1999 – Biodiversité des Ephéméroptères et des Odonates de la Guadeloupe, et biotypologie des cours d'eau de la Basse-Terre. Travail de diplôme, Université de Lausanne/Musée de Zoologie de Lausanne/LEBA (univ. Genève), 60pp + annexes.
- Tsuda, S., 1991 – A distributional list of World Odonata – Osaka, 362 pages.

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|

**NEWLY DISCOVERED ODONATA
LOCALITIES IN THE DOMINICAN
REPUBLIC (WEST INDIES)**

François Meurgey and Carolyne Perron
Museum d'Histoire Naturelle de Nantes (France)
Francois.meurgey@mairie-nantes.fr

Seven Odonata species were found during a collecting trip in the Dominican Republic (Dominican Republic expedition) from 29 January to 7 February 2001. They were all previously known from this island. We list them below as a contribution to Odonata fauna of the Antilles. These specimens stored in the National History Museum of Nantes, under MHNN.Z.0046272 number, and are part of our reference collection for the West Indies. Most of these species were observed at some 100 meters from the right bank of Río San Juan Barracote river, that flows in the North West region of Los Haïtises National Parc and drops to Samana Bay. 63 species are known in Dominican Republic (Daigle, 1993 Bulletin American Odonatology 1 (4) : 65-69 ; Garrison, 1986 Odonatologica 19 (1) : 61-76), but there is a great disparity among the 30 districts ; some have been well surveyed (Distrito Nacional, La Vega and Santiago), but there is no data for 9 of them. Only one species is known for Duarte district to the present (*Lestes forficula*).

ZYGOPTERA

***Telebasis vulnerata* (Hagen, 1861)**

1 m, San Juan Barracote (Los Haïtises), Duarte district, 05.02.2001.

ANISOPTERA

***Erythrodiplax umbrata* L., 1758**

1 f, San Juan Barracote (Los Haïtises), Duarte district, 05.02.2001.

***Scapania frontalis* (Burmeister, 1839)**

1 m, San Francisco de Marcoris, Duarte district, 08.02.2001. Hit by our car on the way to San Francisco de Marconis

***Orthemis ferruginea* (Fabricius, 1775)**

1 f, 1 m, San Juan Barracote (Los Haïtises), Duarte district, 05.02.2001.

***Dythemis rufinervis* (Burmeister, 1839)**

1 f, 1 m, San Juan Barracote (Los Haïtises), Duarte district, 05.02.2001.

***Pantala flavescens* (Fabricius, 1798) :**

1 f, San Juan Barracote (Los Haïtises), Duarte district, 05.02.2001.

***Tramea onusta* Hagen, 1861**

1 m, San Juan Barracote (Los Haïtises), Duarte district, 05.02.2001.

=====

FIRST FRENCH RECORD FOR *ANAX JUNIUS* (DRURY, 1773)

François Meurgey and Carolyne Perron
Museum d'Histoire Naturelle de Nantes (France)
Francois.meurgey@mairie-nantes.fr

Anax junius is a well known and well studied migratory species in North America. The situation is quite different for Europe. *A. junius* has already been observed in Palaearctic region, but only in China and in Kamchatka (Belyshev 1966). The first documented Western Europe observations were made in Autumn 1998, following an early September violent Atlantic storm (remains of hurricane "Earl"). At that time, nine specimens were observed in Cornwall region and on the Scilly Islands, along Great Britain's West coast. No other Western Europe observations related to that species are presently known to us. While studying the migration of *Sympetrum* spp. at Pointe St-Gildas (Loire-Atlantic Department, 02°14 '42.9"W 47°08'09.0"N), an *Anax* specimen was found and first mistaken for the common European *A. imperator*. In order to properly identify this insect, we had to capture the specimen. It is presently stored in the Natural History Museum of Nantes collections. The apex of the four wings indicates that the specimen was old.

This observation, made during the study of the migratory patterns of other species, has interesting implications. American birds or insects, deviating to the East while migrating across Atlantic, usually stop on the first land they find, such as the United Kingdom or Ouessant Island (France), but not generally along the European mainland. The French observation is quite unusual, considering that the length of its Atlantic coast is too great for ease of observation. In early September, only ornithologists are active in this environment (by the way, *A. junius*'s 1998 UK observations were made by ornithologists).

REFERENCES

Balança G., Vissler M.N., 1989.- Observation de la ponte en tandem d'*Anax imperator* Leach, 1815 dans l'Hérault (34) (Odonata, Anisoptera : Aesnidae). Martinia 5 (4) : 90

TRAMEA

The following have been received recently.

"Hey, I created a Dragonfly Game for Environmental Educators. Still need to load an activity map to the site, but the rest is there. < <http://bwwells.org.dragongame>>

Brian Bockhahn, Falls Lake State Park Ranger, Falls & Kerr Lake CBC Compiler, BW Wells Association President, Falls Lake SRA, 13304 Creedmoor Road, Wake Forest NC 27587, 919/676-1027.
<Brian.Bockhahn@ncmail.net>

Kathy Biggs announces her new website, Dragonflies (Odonata) of the Southwest: It showcases the dragons and damsels of the greater southwest, including CA, NV, UT, CO, AZ, & NM.
<http://southwestdragonflies.net>

ARGIA

Binghamton, New York
In This Issue

Vol. 16, No.1, 5 Apr. 2004

		1
Announcing: the 2004 advanced and professional level entomology seminars at the Humboldt Institute on the coast of eastern Maine!		2
Southeast DSA meeting	Carl Cook	2
Updated information		
Other DSA meetings		2
Nymph fest 2004	Michael Veit	3
Aerobatic Anisoptera & zooming Zygoptera: Odonata flight from A to Z. Part 2.	Roy Beckemeyer	4
Dragonfly wings: responding to pressure.		
Skybug		8
Dragonflies of the Northwest Territories, Canada: new records, ecological observations and a checklist	Paul M. Catling, Suzanne Carriere, Deborah Johnson, Mike Fournier	9
Lawsuit filed to establish habitats for dragonfly	Dan Egan	13
Have insects been around since the Silurian?	Roy Beckemeyer	14
Hunt for Red October	Jerrell Daigle	15
First records of pacific spiketail, <i>Cordulegaster dorsalis</i> , in Idaho	Kent Fothergill, John Keebaugh, and Miriam Austin	16
New county records for Iowa - 1999 to 2003	Steve Hummel	18
New Canadian and provincial odonate records for New Brunswick	Dwayne L. Sabine, Anthony W. Thomas and H. Scott Makepeace	22
<i>Anax amazili</i> new for Guadeloupe and notes on other rare species (French West Indies)	François Meurgey and Carolyne Perron	24
Newly discovered odonata localities in the Dominican Republic (West Indies)	François Meurgey and Carolyne Perron	25
First French record for <i>Anax junius</i> (Drury, 1773)	François Meurgey and Carolyne Perron	25
Some Odonata records for the midwest and west for 2003	Roy Beckemeyer	26
<i>Lestes forficula</i> new record from Cayman Islands	Dennis Paulson	26
Dvd review: Common Dragonflies and Damselflies Of Eastern North America, by Dick Walton and Greg Dodge	Allen Barlow	26
Book review: Dragonflies, by Cynthia Berger	Nick Donnelly	27
Book review: Eliza and the Dragonfly, by Susie Caldwell Rinehart	Nick Donnelly	27
Powerpoint review: Dragonfly Dreams, an Album of Summer, by May Lenzer	Nick Donnelly	28
Breaking news- <i>Paltothemis lineatipes</i> explosion in california	Douglas Aguiard	28
TRAMEA		

BACK ISSUES OF ARGIA AND THE BULLETIN OF AMERICAN ODONATOLOGY

The editor is able to provide back issues of **ARGIA**. Please contact T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903. The present price schedule takes into account the different costs of duplication of each number of **ARGIA**. In the event that an issue becomes exhausted, then xerox copies will be sent. **Prices are \$3.00 per issue; these do not include postage; see below.**

Back Issues of the **BULLETIN OF AMERICA ODONATOLOGY** can be furnished at the prices given below. **Prices do not include postage; see below.**

1(1) The Odonata of New York, Thomas W. Donnelly p. 1-28	\$3.00
1(2) Distribution of Dragonflies and Damselflies in Florida, Sidney W. Dunkle p. 29-50	\$2.50
1(3) Morphological and ecological differences among species of <i>Ladona</i> , Michael L. May p. 51-56; Comportamiento reproductivo y policromatismo en <i>Ischnura denticollis</i> Burmeister, Alejandro Córdoba Aguilar [with English summary] p. 57-64	\$1.50
1(4) A checklist of the Odonata of the Dominican Republic by Province, Jerrell James Daigle p. 65-69; Odonata de la Sierra de Huauchinango, Puebla, Mexico [with English summary], José A. Gómez Anaya and Rodolfo Novelo Gutiérrez p. 71-73	\$1.50
2(1) La Náyade de <i>Archilestes latialatus</i> Donnelly, 1981 [with English summary], R. Novelo-Gutiérrez p. 1-7	\$1.50
Descripción e Historia Natural de las Larvas de Odonatos de Costa Rica. III <i>Gynacantha tibiata</i> (Karsch 1891) [with English summary], Alonso Ramírez p. 9-14	
2(2) Description of the Nymph of <i>Epitheca</i> (<i>Tetragoneuria</i>) <i>spinosa</i> (Hagen, K. J. Tennessen p. 15-19	\$1.50
The Larva and Adult Male of <i>Somatochlora georgiana</i> Walker, Jerrell J. Daigle p. 21-26	
2(3) <i>Macromia illinoensis</i> and <i>georgina</i> : a Study of their Variation and Apparent Subspecific Relationship, T.W. Donnelly, K.J. Tennessen p. 27-61	\$3.00
2(4) The Subgenus <i>Tetragoneuria</i> (<i>Anisoptera: Corduliidae: Epitheca</i>) in New Jersey, Michael L. May p. 63-74	\$1.50
3(1) The Odonata of Ohio - a Preliminary Report, Robert C. Glotzhober p. 1-30	\$3.00
3(2) Four Decades of Stability and Change in the Odonata Population at Ten Acre Pond in Central Pennsylvania, Clark N. Shiffer & Harold B. White p. 31-40; Descripción e Historia Natural de las Larvas de Odonatos de Costa Rica. IV. <i>Mecistogaster ornata</i> (Rambur, 1842) [with English summary], Alonso Ramírez p. 43-47	\$1.50
3(3) The Distribution of Odonata in Alabama, Kenneth J. Tennessen, James D. Harper, R. Stephen Krotzer, p. 49-74	\$3.00
3(4) Distribution Records of the Odonata of Montana, Kelly B. Miller and Daniel L. Gustafson, p. 75 - 88	\$1.50
4(1) An Annotated List of the Odonata of New Jersey, With an Appendix on Nomenclature in the Genus <i>Gomphus</i> , Michael L. May & Frank L. Carle p. 1 - 35	\$3.00
4(2) The Odonata of Patuxent Wildlife Research Center and Vicinity, Richard L. Orr p. 37 - 67	\$3.00
4(3) The Status of <i>Lestes apollinaris</i> Navás and <i>L. henschawi</i> Calvert, Thomas W. Donnelly p. 69-74	\$1.50
4(4) The Dragonflies of Washington, Dennis R. Paulson p. 75-90	\$1.50
5(1) The Dragonflies and Damselflies (Odonata) of Louisiana, Bill Mauffray p. 1-26	\$3.00
5(2) The Odonata of the Cayman Islands: a Review, R.R. Askew, R. Prosser, and P.S. Corbet p. 27-32	Taxonomic and \$1.50
Population Studies of British Columbia <i>Aeshna</i> species, G. Peters p. 33-42	
5(3) Adapting the Townes Malaise Trap for Collecting Live Odonata, Robert C. Glotzhober & Dan Riggs, p. 43-48; <i>Archilestes grandis</i> (Great Spreadwing) in Central New Jersey, with Notes on Water Quality, David P. Moskowitz and David M. Bell, p. 49-54; Variation in Head Spines in Female <i>Ophiogomphus</i> , with a Possible Example of Reproductive Character Displacement (Anisoptera: Gomphidae), Dennis R. Paulson, p. 55-58	\$1.50
5(4) The Odonata fauna of Connecticut, David L. Wagner and Michael C. Thomas, p. 59-85	\$3.00
6(1) The Distribution of the Odonata of Hawaii, Jerrell J. Daigle, p. 1-5; Additions to the Description of <i>Gomphomacromia nodisticta</i> Ris 1928 (Odonata: Corduliidae), N. von Ellenrieder, p. 7-11	\$1.50
6(2) The Odonata of Iowa, Robert W. Cruden and O.J. Gode, Jr., p. 13-48	\$3.00
6(3) Odonata in the Great Plains states: Patterns of Distribution and Diversity, Roy J. Beckemeyer, p. 49-99	\$3.00
6(4) Comments on the <i>Erythrodiplax connata</i> (Burmeister, 1839) group, with the elevation of <i>E. fusca</i> (Rambur, 1842), <i>E. minuscula</i> (Rambur, 1842), and <i>E. basifusca</i> (Calvert, 1895) to full species (Anisoptera: Libellulidae), Dennis Paulson, p. 101-110	\$1.50
7(1) The Odonata of the Huron Mountains, Marquette Co., Michigan, M. F. O'Brien, E. Bright & M. A. Kielb, p. 1-22	\$3.00
7(2) Revision of the Order Odonata in Cuba, A. Trapero Q. and C. Naranjo L., p. 23-40	\$1.50
7(3) At-Risk Odonata Of Conterminous United States, George H. Bick pp.41 - 56; Description of the Last Larval Instar of <i>Ischnura fluvialis</i> Selys (Coenagrionidae), Natalia von Ellenrieder and Javier Muzón, 7(3): 57-60	\$1.50
7(4) Distribution of North American Odonata. Part I: Aeshnidae, Petaluridae, Gomphidae, Cordulegastridae, Thomas W. Donnelly, 7(4): 61-90	\$3.00
8(1) Distribution of North American Odonata. Part II: Macromiidae, Corduliidae, and Libellulidae, Thomas W. Donnelly, 8(1): 1-32	*

* subscription

Mailing and Handling Costs:

	SURFACE		AIR MAIL	
	1st number	each additional	1st number	each additional
United States	\$1.25	\$1.00	---	---
Canada, Mexico	\$1.25	\$1.00	\$1.50	\$1.25
Western Hemisphere	\$1.50	\$1.25	\$2.00	\$1.50
Europe, Asia, etc	\$1.50	\$1.25	\$3.00	\$2.50



It may be common, but it sure is pretty! *Pachydiplax longipennis*, by May Lenzer



Everyone's favorite. *Hagenius brevistylus*, by Allen Barlow